arXiv:nlin/0610028v1 [nlin.SI] 12 Oct 2006

The classification of all single travelling wave solutions to Calogero-Degasperis-Focas equation

Chengshi Liu
Department of Mathematics
Daqing Petroleum Institute
Daqing 163318, China
Email: chengshiliu-68@126.com

February 8, 2008

Abstract

Under the travelling wave transformation, Calogero-Degasperis-Focas equation was reduced to an ordinary differential equation. Using a symmetry group of one-parameter, this ODE was reduced to a second order linear inhomogeneous ODE. Furthermore, we applied the change of the variable and complete discrimination system for polynomial to solve the corresponding integrals and obtained the classification of all single travelling wave solutions to Calogero-Degasperis-Focas equation.

Keywords: Classification of travelling wave solution, symmetry group, Calogero-Degasperis-Focas equation

PACS: 02.30.Jr, 05.45.Yv

1 Introduction

In the past decade a lot of expansion methods have been proposed to seek for the travelling wave solutions to nonlinear partial differential equations(for examples see [1-18]). These methods are only indirect methods based on the some assumptions about the forms of the solutions of the equations considered. Applying those indirect methods, we can't give all single travelling wave solutions to the equations considered. On the other hand, some superficially different solutions are essentially the same solution([19]). So it is worthwhile to give the classifications of all single travelling wave solutions to those equations. However, direct integral method is a rather simple and powerful methods([20-23]), if a nonlinear equation can be directly reduced to the integral form as follows:

$$\pm(\xi - \xi_0) = \int \frac{\mathrm{d}u}{\sqrt{p_n(u)}}, \tag{1}$$

where $p_n(u)$ is an n-th order polynomial, we can give the classification([20,23]) of all solutions to the right integral in the Eq.(1) using complete discrimination system for the n-th order polynomial([20-24]).

But there are some nonlinear differential equations whose reduced ODE's are the more complex equations. Therefore we need more technical methods to get the corresponding reduced ODE and its solutions. In the present paper, using the trick of symmetry group to reduce the Calogero-Degasperis-Focas equation([25]) to an integrable ODE, and furthermore using the change of variable and complete discrimination system for polynomial to solve the corresponding integral, we obtain the classification of all single travelling wave solutions to Calogero-Degasperis-Focas equation. The method and trick used here have general meaning for the studies of travelling wave solutions to some nonlinear partial differential equations.

2 Classification of all single travelling wave solutions to Calogero-Degasperis-Focas equation

Calogero-Degasperis-Focas equation reads([25])

$$u_t = u_{xxx} - \frac{1}{8}u_x^3 + (a\exp u + b\exp(-u) + c)u_x. \tag{2}$$

Take the travelling wave transformation $u = u(\xi), \quad \xi = kx + \omega t$. The corresponding reduced ODE is as follows:

$$\omega u' = k^3 u''' - \frac{k^3}{8}(u')^3 + (a\exp u + b\exp(-u) + c)ku'. \tag{3}$$

Since the Eq.(2) is invariant under the following one-parameter group

$$u* = u, \tag{4}$$

$$\xi* = \xi + \varepsilon, \tag{5}$$

so we can reduce the order of the Eq.(3) by one. For the purpose, we take $z = u'$, then we have $u'' = z\frac{\mathrm{d}z}{\mathrm{d}u}$ and $u''' = z(\frac{\mathrm{d}z}{\mathrm{d}u})^2 + z^2\frac{\mathrm{d}^2z}{\mathrm{d}u^2}$. Instituting those terms into the Eq.(3) and furthermore letting $W = z^2$, we have

$$W'' - \frac{1}{4}W = \frac{2\omega}{k^3} - \frac{2}{k^2}(a\exp u + b\exp(-u) + c). \tag{6}$$

Using the method of the variation of constants, the general solution of the Eq.(6) is as follows:

$$W(u) = c_1\exp(\frac{1}{2}u) + c_2\exp(-\frac{1}{2}u) - \frac{8a}{3k^2}\exp(u) - \frac{8b}{3k^2}\exp(-u) - 8(\frac{\omega}{k^3} - \frac{c}{k^2}), \tag{7}$$

where c_1 and c_2 are two arbitrary constants. Hence we have the following result:

Theorem: the general solution of the Eq.(3) is as follows:

$$\pm(\xi-\xi_0)=\int\frac{\mathrm{d}u}{\sqrt{c_1\exp(\frac{1}{2}u)+c_2\exp(-\frac{1}{2}u)-\frac{8a}{3k^2}\exp(u)-\frac{8b}{3k^2}\exp(-u)-8(\frac{\omega}{k^3}-\frac{c}{k^2})}}. \quad (8)$$

In order to solve the integral (8),we take a variable transformation $v=\exp(\frac{1}{2}u)$, the corresponding integral becomes

$$\pm\frac{1}{2}(\xi-\xi_0)=\int\frac{\mathrm{d}v}{\sqrt{a_4v^4+a_3v^3+a_2v^2+a_1v+a_0}}. \quad (9)$$

where

$$a_4=-\frac{8a}{3k^2},\quad a_3=c_1,\quad a_2=-\frac{8}{k^2}(\frac{\omega}{k}-c),\quad a_1=c_2,\quad a_0=-\frac{8b}{3k^2}. \quad (10)$$

If $a<0$, we take the change of variable as follows:

$$w=a_4^{\frac{1}{4}}(v+\frac{a_3}{4a_4}), \quad (11)$$

Then Eq.(9) becomes

$$\pm\frac{1}{2}a_4^{\frac{1}{4}}(\xi-\xi_0)=\int\frac{\mathrm{d}w}{\sqrt{w^4+pw^2+qu+r}}, \quad (12)$$

where

$$p=\frac{a_2}{\sqrt{a_4}},\quad q=a_4^{-\frac{1}{4}}(\frac{a_3^3}{8a_4^2}-\frac{a_2a_3}{2a_4}+a_1),\quad r=a_0-\frac{a_1a_3}{4a_4}+\frac{a_2a_3^2}{16a_4^2}-\frac{3a_3^4}{256a_4^3}. \quad (13)$$

If $a>0$, we take the change of variable as follows:

$$w=(-a_4)^{\frac{1}{4}}(v+\frac{a_3}{4a_4}), \quad (14)$$

Then Eq.(9) becomes

$$\pm\frac{1}{2}(-a_4)^{\frac{1}{4}}(\xi-\xi_0)=\int\frac{\mathrm{d}w}{\sqrt{-(w^4+pw^2+qu+r)}}, \quad (15)$$

where

$$p=-\frac{a_2}{\sqrt{-a_4}},\quad q=(-a_4)^{-\frac{1}{4}}(-\frac{a_3^3}{8a_4^2}+\frac{a_2a_3}{2a_4}-a_1),\quad r=-a_0-\frac{a_1a_3}{4a_4}-\frac{a_2a_3^2}{16a_4^2}+\frac{3a_3^4}{256a_4^3}. \quad (16)$$

According to the classification([23]) of the solutions to the ODE $(w')^2 = \epsilon(w^4 + pw^2 + qw + r)$, where $\epsilon = \pm 1$, we can give the classification of all single travelling wave solutions to Calogero-Degasperis-Focas equation. We denote

$$f(w) = w^4 + pw^2 + qw + r, \tag{17}$$

and write its complete discrimination system as follows([23]):

$$D_1 = 4, D_2 = -p, D_3 = 8rp - 2p^3 - 9q^2,$$
$$D_4 = 4p^4 r - p^3 q^2 + 36prq^2 - 32r^2 p^2 - \frac{27}{4}q^4 + 64r^3,$$

$$F_2 = 9p^2 - 32pr. \tag{18}$$

According to the complete discrimination system (18) and the above theorem, we obtain the classification of all single travelling wave solutions to Calogero-Degasperis-Focas equation.

Case 1: If $D_4 = 0, D_3 = 0, D_2 < 0$, then we have

$$f(w) = ((w - l_1)^2 + s_1^2)^2, \tag{19}$$

where l_1, s_1 are real numbers, $s_1 > 0$. When $\epsilon = +1$, we have

$$u = 2\ln\{a_4^{\frac{1}{4}}[s_1 \tan(\frac{1}{2}a_4^{\frac{1}{4}} s_1(\xi - \xi_0)) + l_1] - \frac{a_3}{4a_4}\}. \tag{20}$$

Case 2: If $D_4 = 0, D_3 = 0, D_2 = 0$, then we have

$$f(w) = w^4. \tag{21}$$

When $\epsilon = 1$, we have

$$u = 2\ln[\pm\frac{2a_4^{-\frac{1}{2}}}{\xi - \xi_0} - \frac{a_3}{4a_4}]. \tag{22}$$

Case 3: If $D_4 = 0, D_3 = 0, D_2 > 0, E_2 > 0$, then we have

$$f(w) = (w - \alpha)^2 (w - \beta)^2, \tag{23}$$

where α, β, are real numbers, $\alpha > \beta$. If $\epsilon = 1$, when $w > \alpha$ or $w < \beta$, we have

$$u = 2\ln\{a_4^{-\frac{1}{4}}[\frac{\beta - \alpha}{2}(\coth\frac{\alpha - \beta}{4}a_4^{\frac{1}{4}}(\xi - \xi_0) - 1) + \beta] - \frac{a_3}{4a_4}\}; \tag{24}$$

when $\alpha > w > \beta$, we have

$$u = 2\ln\{a_4^{-\frac{1}{4}}[\frac{\beta-\alpha}{2}(\tanh\frac{\alpha-\beta}{4}a_4^{\frac{1}{4}}(\xi-\xi_0)-1)+\beta]-\frac{a_3}{4a_4}\}. \tag{25}$$

Case 4: If $D_4 = 0, D_3 > 0, D_2 > 0$, then we have

$$f(w) = (w-\alpha)^2(w-\beta)(w-\gamma), \tag{26}$$

where α, β, γ are real numbers, $\beta > \gamma$. If $\epsilon = 1$, when $\alpha > \beta$ and $w > \beta$, or when $\alpha < \gamma$ and $w < \gamma$, we have

$$\exp\{\pm\frac{a_4^{\frac{1}{4}}\sqrt{(\alpha-\beta)(\alpha-\gamma)}(\xi-\xi_0)}{2}\} = \frac{\{\sqrt{[a_4^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\beta](\alpha-\gamma)}-\sqrt{(\alpha-\beta)[a_4^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\gamma]}\}^2}{|a_4^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\alpha|}; \tag{27}$$

when $\alpha > \beta$ and $w < \gamma$, or when $\alpha < \gamma$ and $w < \beta$, we have

$$\exp\{\pm\frac{a_4^{\frac{1}{4}}\sqrt{(\alpha-\beta)(\alpha-\gamma)}(\xi-\xi_0)}{2}\} = \frac{\{\sqrt{[a_4^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\beta](\gamma-\alpha)}-\sqrt{(\beta-\alpha)[a_4^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\gamma]}\}^2}{|a_4^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\alpha|}; \tag{28}$$

when $\beta > \alpha > \gamma$, we have

$$\pm\sin\{\frac{a_4^{\frac{1}{4}}\sqrt{(\beta-\alpha)(\alpha-\gamma)}(\xi-\xi_0)}{2}\} = \frac{[a_4^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\beta](\alpha-\gamma)+(\alpha-\beta)[a_4^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\gamma]}{|[a_4^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\alpha](\beta-\gamma)|}. \tag{29}$$

If $\epsilon = -1$, when $\alpha > \beta$ and $w > \beta$, or when $\alpha < \gamma$ and $w < \gamma$, we have

$$\exp\{\pm\frac{(-a_4)^{\frac{1}{4}}\sqrt{(\alpha-\beta)(\alpha-\gamma)}(\xi-\xi_0)}{2}\} = \frac{\{\sqrt{[(-a_4)^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\beta](\alpha-\gamma)}-\sqrt{(\alpha-\beta)[(-a_4)^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\gamma]}\}^2}{|(-a_4)^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\alpha|}; \tag{30}$$

when $\alpha > \beta$ and $w < \gamma$, or When $\alpha < \gamma$ and $w < \beta$, we have

$$\exp\{\pm\frac{(-a_4)^{\frac{1}{4}}\sqrt{(\alpha-\beta)(\alpha-\gamma)}(\xi-\xi_0)}{2}\} =$$

$$\frac{\{\sqrt{[(-a_4)^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\beta](\gamma-\alpha)}-\sqrt{(\beta-\alpha)[(-a_4)^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\gamma]}\}^2}{|(-a_4)^{\frac{1}{4}}[\exp(\frac{u}{2})+\frac{a_3}{4a_4}]-\alpha|}; \quad (31)$$

when $\beta > \alpha > \gamma$, we have

$$\pm\sin\{\frac{(-a_4)^{\frac{1}{4}}\sqrt{(\beta-\alpha)(\alpha-\gamma)}(\xi-\xi_0)}{2}\} = \frac{[(-a_4)^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\beta](\alpha-\gamma)+(\alpha-\beta)[(-a_4)^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\gamma]}{|[(-a_4)^{\frac{1}{4}}(\exp(\frac{u}{2})+\frac{a_3}{4a_4})-\alpha](\beta-\gamma)|}. \quad (32)$$

Case 5: If $D_4 = 0, D_3 = 0, D_2 > 0, E_2 = 0$, then we have

$$f(w) = (w-\alpha)^3(w-\beta), \quad (33)$$

where α, β are real numbers. If $\epsilon = 1$, when $w > \alpha, w > \beta$; or $w < \alpha, w < \beta$, We have

$$u = 2\ln\{a_4^{-\frac{1}{4}}[\frac{4(\alpha-\beta)}{\frac{1}{4}a_4^{\frac{1}{2}}(\alpha-\beta)^2(\xi-\xi_0)^2-4}+\alpha]-\frac{a_3}{4a_4}\}. \quad (34)$$

If $\epsilon = -1$, when $w > \alpha, w < \beta$; or $w < \alpha, w > \beta$, we have

$$u = 2\ln\{(-a_4)^{-\frac{1}{4}}[\frac{4(\beta-\alpha)}{4+\frac{1}{4}(-a_4)^{\frac{1}{2}}(\alpha-\beta)^2(\xi-\xi_0)^2}+\alpha]-\frac{a_3}{4a_4}\}. \quad (35)$$

Case 6: If $D_4 = 0, D_2D_3 < 0$, then we have

$$f(w) = (w-\alpha)^2((w-l_1)^2+s_1^2), \quad (36)$$

where α, l_1 and s_1 are real numbers. If $\epsilon = 1$, we have

$$u = 2\ln\{a_4^{-\frac{1}{4}} \times [\frac{\exp[\pm\frac{1}{2}a_4^{\frac{1}{4}}\sqrt{(\alpha-l_1)^2+s_1^2}(\xi-\xi_0)]-\gamma+(2-\gamma)\sqrt{(\alpha-l_1)^2+s_1^2}}{\{\exp[\pm\frac{1}{2}a_4^{\frac{1}{4}}\sqrt{(\alpha-l_1)^2+s_1^2}(\xi-\xi_0)]-\gamma\}^2-1}] - \frac{a_3}{4a_4}\}. \quad (37)$$

where

$$\gamma = \frac{\alpha-2l_1}{\sqrt{(\alpha-l_1)^2+s_1^2}}. \quad (38)$$

Case 7: If $D_4 > 0, D_3 > 0, D_2 > 0$, then we have

$$f(w) = (w-\alpha_1)(w-\alpha_2)(w-\alpha_3)(w-\alpha_4), \quad (39)$$

where $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ are real numbers. and $\alpha_1 > \alpha_2 > \alpha_3 > \alpha_4$. If $\epsilon = 1$, when $W > \alpha_1$ or $w < \alpha_4$, we have

$$u = 2\ln\{a_4^{-\frac{1}{4}} \times [\frac{\alpha_2(\alpha_1-\alpha_4)\,\mathrm{sn}^2\,(\frac{a_4^{\frac{1}{4}}\sqrt{(\alpha_1-\alpha_3)(\alpha_2-\alpha_4)}}{4}(\xi-\xi_0), m) - \alpha_1(\alpha_2-\alpha_4)}{(\alpha_1-\alpha_4)\,\mathrm{sn}^2\,(\frac{a_4^{\frac{1}{4}}\sqrt{(\alpha_1-\alpha_3)(\alpha_2-\alpha_4)}}{4}(\xi-\xi_0), m) - (\alpha_2-\alpha_4)}] - \frac{a_3}{4a_4}\}; \tag{40}$$

when $\alpha_2 > w > \alpha_3$, we have

$$u = 2\ln\{a_4^{-\frac{1}{4}} \times [\frac{\alpha_4(\alpha_2-\alpha_3)\,\mathrm{sn}^2\,(\frac{a_4^{\frac{1}{4}}\sqrt{(\alpha_1-\alpha_3)(\alpha_2-\alpha_4)}}{4}(\xi-\xi_0), m) - \alpha_3(\alpha_2-\alpha_4)}{(\alpha_2-\alpha_3)\,\mathrm{sn}^2\,(\frac{a_4^{\frac{1}{4}}\sqrt{(\alpha_1-\alpha_3)(\alpha_2-\alpha_4)}}{4}(\xi-\xi_0), m) - (\alpha_2-\alpha_4)}] - \frac{a_3}{4a_4}\}. \tag{41}$$

If $\epsilon = -1$, when $\alpha_1 > w > \alpha_2$, we have

$$u = 2\ln\{(-a_4)^{-\frac{1}{4}} \times [\frac{\alpha_3(\alpha_1-\alpha_2)\,\mathrm{sn}^2\,(\frac{(-a_4)^{\frac{1}{4}}\sqrt{(\alpha_1-\alpha_3)(\alpha_2-\alpha_4)}}{4}(\xi-\xi_0), m) - \alpha_2(\alpha_1-\alpha_3)}{(\alpha_1-\alpha_2)\,\mathrm{sn}^2\,(\frac{(-a_4)^{\frac{1}{4}}\sqrt{(\alpha_1-\alpha_3)(\alpha_2-\alpha_4)}}{4}(\xi-\xi_0), m) - (\alpha_1-\alpha_3)}] - \frac{a_3}{4a_4}\}; \tag{42}$$

when $\alpha_3 > w > \alpha_4$, we have

$$u = 2\ln\{(-a_4)^{-\frac{1}{4}} \times [\frac{\alpha_1(\alpha_3-\alpha_4)\,\mathrm{sn}^2\,(\frac{(-a_4)^{\frac{1}{4}}\sqrt{(\alpha_1-\alpha_3)(\alpha_2-\alpha_4)}}{4}(\xi-\xi_0), m) - \alpha_4(\alpha_3-\alpha_1)}{(\alpha_3-\alpha_4)\,\mathrm{sn}^2\,(\frac{(-a_4)^{\frac{1}{4}}\sqrt{(\alpha_1-\alpha_3)(\alpha_2-\alpha_4)}}{4}(\xi-\xi_0), m) - (\alpha_3-\alpha_1)}] - \frac{a_3}{4a_4}\}, \tag{43}$$

where $m^2 = \frac{(\alpha_1-\alpha_2)(\alpha_3-\alpha_4)}{(\alpha_1-\alpha_3)(\alpha_2-\alpha_4)}$.

Case 8: If $D_4 < 0, D_2D_3 \geq 0$, then we have

$$f(w) = (w-\alpha)(w-\beta)((w-l_1)^2 + s_1^2), \tag{44}$$

where α, β, l_1 and s_1 are real numbers, and $\alpha > \beta, s_1 > 0$, we have

$$u = 2\ln\{(\epsilon a_4)^{-\frac{1}{4}}[\frac{a\,\mathrm{cn}\,(\frac{(\epsilon a_4)^{\frac{1}{4}}\sqrt{-\epsilon 2 s_1 m_1(\alpha-\beta)}}{4mm_1}(\xi-\xi_0), m) + b}{c\,\mathrm{cn}\,(\frac{(\epsilon a_4)^{\frac{1}{4}}\sqrt{-\epsilon 2 s_1 m_1(\alpha-\beta)}}{4mm_1}(\xi-\xi_0), m) + d}] - \frac{a_3}{4a_4}\}. \tag{45}$$

where

$$c=\alpha-l_1-\frac{s_1}{m_1}, d=\alpha-l_1-s_1m_1,$$

$$a=\frac{1}{2}[(\alpha+\beta)c-(\alpha-\beta)d], b=\frac{1}{2}[(\alpha+\beta)d-(\alpha-\beta)c],$$

$$E=\frac{s_1^2+(\alpha-l_1)(\beta-l_1)}{s_1(\alpha-\beta)},\quad m_1=E\pm\sqrt{E^2+1}, m^2=\frac{1}{1+m_1^2}, \tag{46}$$

we choose m_1 such that $\epsilon m_1<0$.

Case 9: If $D_4>0, D_2D_3\leq 0$, then we have

$$f(w)=((w-l_1)^2+s_1^2)(w-l_2)^2+s_2^2), \tag{47}$$

where l_1, l_2, s_1 and s_2 are real numbers, and $s_1>s_2>0$. If $\epsilon=1$, we have

$$u=2\ln\{a_4^{-\frac{1}{4}}[\frac{a\,\mathrm{sn}\,(\eta(\xi-\xi_0),m)+b\,\mathrm{cn}\,(\eta(\xi-\xi_0),m)}{c\,\mathrm{sn}\,(\eta(\xi-\xi_0),m)+d\,\mathrm{cn}\,(\eta(\xi-\xi_0),m)}]-\frac{a_3}{4a_4}\}, \tag{48}$$

where

$$c=-s_1-\frac{s_2}{m_1},\quad d=l_1-l_2,\quad a=l_1c+s_1d,\quad b=l_1d-s_1c,$$

$$E=\frac{s_1^2+s_2^2+(l_1-l_2)^2}{2s_1s_2},\quad m_1=E+\sqrt{E^2-1},$$

$$m^2=1-\frac{1}{m_1^2},\quad \eta=\frac{1}{2}a_4^{\frac{1}{4}}s_2\sqrt{\frac{m_1^2c^2+d^2}{c^2+d^2}}. \tag{49}$$

These are all possible cases, so we obtain the complete classification of all single travelling wave solutions to Calogero-Degasperis-Focas equation.

3 Conclusions

In summary, although the classifications of all single travelling wave solutions to nonlinear partial differential equations is a rather difficult problem. But there are a lot of nonlinear differential equations whose all travelling wave solutions can be obtained using direct integral method and complete discrimination system for polynomial. On the other hand, if a nonlinear differential equation whose reduced ODE can't be obtained by simple integral method, then we will need find more powerful tricks and methods to do this thing. In this paper, we use symmetry group to reduce the order of ODE, and furthermore reduce the equation to an integrable ODE. Using complete discrimination system for polynomial, we obtain the classifications of all single travelling wave solutions to some nonlinear partial differential equations. The methods and tricks used here can be expected to develop to solve more complex and more extensive equations.

References

[1] Wang M L,Zhou Y B.,Li Z B. Phy. Lett. 1996; A216: 67

[2] Wang M L Phys. Lett. 1995; A199: 169

[3] Fan E G Phys.Lett . 2000; A277:212.

[4] Liu S K, Fu Z T, Liu S D, Zhao Q Phys. Lett. A 2001; A290: 72.

[5] Yan Z Y ,Chaos,Solitons and Fractals. 2003;15 :575

[6] Lou S Y, Lu J Z, J. Phys. A. 1996; 29: 4029

[7] Fan E G J. Phys. Soc. Japan. 2002; 71: 2663.

[8] Fan E G Chaos, Solitons and Fractals. 2003;16: 819

[9] Fan E G J. Phys. A, 2002 35 6853

[10] Fan E G J. Phys. A, 2000 33 6925

[11] Liu C S Acta. Phys. Sin. 2005; 54. 2505(in Chinese)

[12] Malfliet W Am. J. Phys. 1992 60 650

[13] Parkes E J and Duffy B R Comput.Phys.Commu.1996 98 288

[14] Elwakil S A et al,Phys.Lett. A. 2002 299 179

[15] Yan Z Y Phys.Lett. A. 2001 285 355

[16] Li B et al J. Phys. A, 2003

[17] Conte R and Mustte M J.Phys.A.1992 25 5609

[18] Bountis T C et al J.Math,Phys.1986 27n1215

[19] Liu C P, Commu. Theor. Phys. 2005 44 609

[20] Liu C S Chin. Phys. Lett. 2004; 21:2369.

[21] Liu C S and Du X H Acta. Phys. Sin. 2005; 54: 1039. (in Chinese)

[22] Liu C S Commun.Thero. Phys. 2005; 43: 787

[23] Liu C S Chin. Phys. 2005 14 1710

[24] Yang L, Hou X R, Zeng Z B Sci. in China, 1996; Ser.E 39: 628.

[25] Ablowicz M J, Kruskal M D and Ladik J F SIAM J Appl Math. 1979 36 428